AUTHOR INDEX, VOLUME 6

Alexandrowicz, Z., and Daniel, E.: On the limiting sedimentation coefficients of polyelectrolytes, 1500
Applequist, J.: Comments on the one-dimensional Ising model and its application to the *cis–trans* equilibrium in poly-L-proline, 117
Augustat, S.: see Szejtli, J.

Bach, D.: see Miller, I. R.
Baer, E.: see Carr, S. H.
Balazs, E. A.: see Gibbs, D. A.
Bangerter, B. W., and Chan, S. I.: Proton magnetic resonance study of the binding of purine to polyuridylic acid, 983
Berger, A.: see Ingwall, R. T.
Birdi, K. S.: see Hoyer, H. W.
Black, D. R.: see Kasarda, D. D.
Blake, A., and Peacocke, A. R.: The interaction of aminoacridines with nucleic acids, 1225
Bloomfield, V. A.: Sedimentation coefficients of replicating and crosslinked DNA, 285
—: see Douthart, R. J.; Sharp, P.
Blout, E. R.: see Morita, K.
Boni, R.: see Giacometti, G.
Bradbury, E. M., Carpenter, B. G., and Goldman, H.: Conformational studies of polymers and copolymers of L-aspartate esters. I. Preparation and solution studies, 837
—, Carpenter, B. G., and Stephens, R. M.: Conformational studies of polymers and copolymers of L-aspartate esters. II. Infrared studies and the factors involved in the formation of the ω-helix, 905
—, Crane-Robinson, C., Goldman, H., and Rattle, H. W. E.: Proton magnetic resonance and optical spectroscopic studies of water-soluble polypeptides: Poly-L-lysine HBr, poly(L-glutamic acid), and copoly(L-glutamic acid^{42}, L-lysine HBr^{28}, L-alanine^{30}), 851
Breuer, M. M.: Strain–stress isotherm of keratin fibers, 1503
Bumb, R. R.: see Jackobs, J. J.
Burke, J. J.: see Threlkeld, J. O.

Cantor, C. R.: Kinetics of primer-dependent polynucleotide phosphorylase synthetic reactions, 369
—, and Chin, W. W.: Oligonucleotide interactions. I. Structure of 2:1 complexes between polyuridylic acid and oligoadenylic acid, 1745
Carmichael, J. B.: Theory of gel filtration separation of biopolymers assuming a Gaussian distribution of pore sizes, 1497
Carpenter, B. G.: see Bradbury, E. M.
Carr, S. H., Walton, A. G., and Baer, E.: Epitaxial crystallization of poly-(γ-benzyl L-glutamate on alkali halide single crystals, 469
Casimir, W. v., Kaiser, N., Keilmann, F., Mayer, A., and Vogel, H.: Dielectric properties of oxyhemoglobin and deoxyhemoglobin in aqueous solution at microwave frequencies, 1705
Chakravarti, K.: see Chowrashi, P. K.
Chan, S. I.: see Bangerter, B. W.
Chandrasekharan, R.: see Ramachandran, G. N.
Chattoraj, D. K.: see Chowrashi, P. K.
Chin, W. W.: see Cantor, C. R.
Chowrashi, P. K., Chattoraj, D. K., and Chakravarti, K.: Electrophoresis of absorbed RNA, 97
Ciferri, A., Puett, D., Rajagh, L., and Hermans, J., Jr.: Potentiometric titrations and the helix–coil transition of poly(L-glutamic acid) and poly-L-lysine in aqueous salt solutions, 1019
—: see Puett, D.; Threlkeld, J. O.
Cleland, R. L.: Ionic polysaccharides. II. Comparison of polyelectrolyte behavior of hyaluronate with that of carboxymethyl cellulose, 1519
Cohen, G., and Eisenberg, H.: Deoxyribonucleate solutions: Sedimentation in a density gradient, partial specific volumes, density and refractive index increments, and preferential interactions, 1077
Constantin, D.: see Spach, G.

Cosani, A., Peggion, E., Verdini, A. S., and Terbojevich, M.: Far ultraviolet optical rotatory properties of poly-L-tryptophan. I, 963
—: see Peggion, E.
Craig, M. E., and Crothers, D. M.: Calculation of kinetic curves for the helix–coil transition of polypeptides, 385
Crane-Robinson, C.: see Bradbury, E. M.
Crothers, D. M.: Calculation of binding isotherms for heterogeneous polymers, 575. Calculation of melting curves for DNA, 1391
—: see Craig, M. E.; Fink, T. R.

Daniel, E.: see Alexandrowicz, Z.
Davidson, N.: see Lee, C. S.; Yew, F. F. H.
Davis, R. C., and Tinoco, I., Jr.: Temperature-dependent properties of dinucleoside phosphates, 223
Del Pra, A.: see Peggion, E.
De Santis, P., and Kovacs, A. J.: Molecular conformation of poly(S-lactic acid), 299
Douthart, R. J., and Bloomfield, V. A.: Intrinsic viscosity of cyclic and linear λ DNA, 1297

Eisenberg, H.: see Cohen, G.
Elson, E.: Kinetics of synthesis of polymeric dAT on oligomeric templates, 269
Epand, R. F., and Scheraga, H. A.: The helix–coil transition of poly-L-lysine in methanol–water solvent mixtures 1383. Conformations of poly-L-valine in solution, 1551
Eyring, E. J., Kraus, H., and Yang, J. T.: Induced optical activity of chondroitin sulfate C–acridine orange complexes, 703
—, and Yang, J. T.: Viscosity and optical activity of chondroitin sulfate C, 691

Falxa, M. L.: see Toniolo, C.
Fasman, G. D.: see Lynn, J.; Tooney, N. M.
Feughelman, M., and Mitchell, T. W.: The state of β-keratin in thermally denatured keratin fibers, 1515
Fink, T. R., and Crothers, D. M.: Comparison of several calculations of helix–coil transitions in heterogeneous polymers, 863
Flemming, J.: Über die Adsorption von DNS in der Grenzfläche Quecksilber–Elektrolyt, 1697
Frič, J.: see Zimmer, C.
Frommer, M. A., and Miller, I. R.: Influence of the tritium labeling of calf thymus DNA by the Wilzbach method on its physical properties, 1461
Fujishiro, R.: see Kagemoto, A.
Fukutome, H.: see Yamada, T.

Gabbay, E. J., Kleinman, R., and Shimshak, R. R.: Topography of nucleic acid helices in solution. VI. The effect of amino acid derivatives on the RNAse-catalyzed hydrolysis of polyadenylic acid. Demonstration of an asymmetric surface, 993
—, and Shimshak, R. R.: Topography of nucleic acid helices in solutions. IV. Effect of polyamines on RNAse-catalyzed hydrolysis of polyadenylic acid, 255
—: see Glaser, R.
Gabor, G.: Radiationless energy transfer through a polypeptide chain, 809
Galley, W. C.: On the triplet states of polynucleotide–acridine complexes. I. Triplet energy delocalization in the 9-aminoacridine–DNA complex, 1279
Gennes, P.-G. de: Statistics of branching and hairpin helices for the dAT copolymer, 715
Ghatak, U. R.: see Kovacs, J.
Giacometti, G., Turolla, A., and Boni, R.: Enthalpy of helix–coil transitions from heats of solution. I. Polyglutamates, 441
Gibbs, D. A., Merrill, E. W., Smith, K. A., and Balazs, E. A.: Rheology of hyaluronic acid, 777
Gibbs, J. H.: see MacDonald, C. T.
Glaser, R., and Gabbay, E. J.: Topography of nucleic acid helices in solutions. III. Interactions of spermine and spermidine derivatives with polyadenylic–polyuridylic and polyinosinic–polycytidylic acid helices, 243
Glaubiger, D., Lloyd, D. A., and Tinoco, I., Jr.: Temperature-dependent optical properties of a torsional oscillator model for dinucleoside phosphates, 409
Goel, N. S.: Relaxation kinetics of denaturation of DNA, 55

—, and Montroll, E. W.: Denaturation and renaturation of DNA. II. Possible use of synthetic periodic copolymers to establish model and parameters, 731
Goodman, M., and Toniolo, C.: Conformational studies on proteins with aromatic side-chain effects, 1673
—, —, and Peggion, E.: Conformational aspects of polypeptides. XXIX. Conformational assignments for some aromatic polypeptides by far-UV cotton effects. New results, 1691
—: see Toniolo, C.
Goldman, H.: see Bradbury, E. M.
Goldsack, D. E.: Volume fraction of the polypeptide chain in proteins and the concentration of protein denaturants, 164
Graves, I. L.: Interaction of heat-denatured HeLa cell DNA with synthetic and natural polysaccharides, 1573
Gray, D. M., and Rubenstein, I.: Ultraviolet dichroic ratio of DNA from T2 and T5 bacteriophages, 1605
Green, G., and Mahler, H. R.: Optical properties of DNA in ethylene glycol, 1509

Haly, A. R., and Snaith, J. W.: Specific heat studies of various wool–water systems, 1355
Harpst, J. A., Krasna, A. I., and Zimm, B. H.: Low-angle light-scattering instrument for DNA solutions, 585. Molecular weight of T7 and calf thymus DNA by low-angle light scattering, 595
Harrington, R. E.: Flow birefringence of T2 bacteriophage DNA, 105
Hayashi, T.: see Nakajima, A.
Hearst, J. E.: see Tunis, M.-J. B.
Hermans, J., Jr.: see Ciferri, A.
Hinkle, M. E., and Zobel, H. F.: X-Ray diffraction of oriented amylose fibers. III. The structure of amylose–*n*-butanol complexes, 1119
Holcomb, D. N., and Timasheff, S. N.: Temperature dependence of the hydrogen ion equilibria in poly(riboadenylic acid), 513
Hoyer, H. W., and Birdi, K. S.: Detection and enthalpy of vaporization of bound water in biopolymers by DTA, 1507

Ikegami, A.: Hydration of polyacids, 431
Ingwall, R. T., Scheraga, H. A., Lotan, N., Berger, A., and Katchalski, E.: Conformational studies of poly-L-alanine in water, 331
Isemura, T., Okabayashi, H., and Sakakibara, S.: Steric structure of L-proline oligopeptides. I. Infrared absorption spectra of the oligopeptides and poly-L-proline, 307
—: see Okabayashi, H.
Itoh, K., Nakahara, T., Shimanouchi, T., Oya, M., Uno, K., and Iwakura, Y.: Far-infrared spectra of polyalanines with α-helical and β-form structures, 1759
Iwakura, Y.: see Itoh, K.

Jackobs, J. J., Bumb, R. R., and Zaslow, B.: Crystalline structure in oriented fibers of KBr–amylose, 1659
Jennings, B. R.: An estimation of the dimensions of the phycocyanin molecule, 1177
—, Spach, G., and Schuster, T. M.: Specific aggregation of poly(α-L-glutamic acid) and hysteresis effects in aqueous solutions. I. Influence of temperature-dependent aggregation on the optical rotation of PGA, 635
Jones, D. W.: A minimization approach for the improvement of cellulose structure models, 771

Kagemoto, A., and Fujishiro, R.: Ethalpy change of the coil–helix transition of poly(γ-benzyl L-glutamate) in dichloroacetic acid–1,2-dichloroethane mixtures, 1753
Kaiser, N.: see Casimir, W. v.
Kasarda, D. D., and Black, D. R.: Thermal degradation of proteins studied by mass spectrometry, 1001
Katchalski, E.: see Ingwall, R. T.
Katchalsky, A.: see Reich, S.
Keilmann, F.: see Casimir, W. v.
Kelly, R. B.: see Strauss, J. H., Jr.
Kleinman, R.: see Gabbay, E. J.
Klotz, I. M., and Lyndrup, M. L.: Acid–base behavior of poly(vinylpyrrolidone–vinylimidazole) copolymers 1405
Kobayasi, S.: Decay processes of the electric birefringence in polydispersed systems, 1491

Kopple, K. D.: Esterolytic activity of vinylimidazole–vinylpyrrolidinone copolymers, 1417
Kotýnek, O.: see Zikán, J.
Kovacs, A. J.: see De Santis, P.
Kovacs, J., Schmit, G. N., and Ghatak, U. R.: Polypeptides with known repeating sequences of amino acids. Comparison of several methods used for the synthesis of poly-γ-D- and L-glutamylglycine and investigation of its serological reaction with antianthrax immune serum, 817
Koyama, Y., and Shimanouchi, T.: Conformations of model compounds of proteins. I. Acetylglycine *N*-methylamide, 1037
Krakauer, H., and Sturtevant, J. M.: Heats of the helix–coil transitions of the poly A–poly U complexes, 491
Krasna, A. I.: see Harpst, J. A.
Kraus, H.: see Eyring, E. J.
Krimm, S., and Kuroiwa, K.: Low temperature infrared spectra of polyglycines and C—H···O=C hydrogen bonding in polyglycine II, 401
—: see Tiffany, M. L.
Kubota, S., Sugai, S., and Noguchi, J.: Conformations and optical rotatory properties of poly-*O*-acetylthreonine and poly-*O*-acetylallothreonine, 1311
Kuroiwa, K.: see Krimm, S.

Lee, C. S., and Davidson, N.: Flow dichroism of deoxyribonucleic acid solutions, 531
Lloyd, D. A.: see Glaubiger, D.
Lotan, N.: see Ingwall, R. T.
Luck, G.: see Zimmer, C.
Lyndrup, M. L.: see Klotz, I. M.
Lynn, J., and Fasman, G. D.: Conformational dependence of fluorescence polarization spectra of L-tryptophan containing copolypeptides, 159

MacDonald, C. T., Gibbs, J. H., and Pipkin, A. C.: Kinetics of biopolymerization on nucleic acid templates, 1
McMenamy, R. H.: see Pande, C. S.
Maestre, M. F.: Transient electric birefringence studies of T2 bacteriophage and T2 ghost, 415
Mahler, H. R.: see Green, G.
Makino, S., Wakabayashi, K., and Sugai, S.: Ion-binding and conformation of poly-L-methionone *S*-methylsulfonium salts in added salt systems, 551
Mammi, M.: see Peggion, E.
Masuzawa, M., and Sterling, C.: Gel–water relationships: Dielectric dispersion, 1453
Matsumoto, M., Watanabe, H., and Yoshioka, K.: Electrical and hydrodynamical properties of polypeptides in solution. I. Poly(L-glutamic acid) in in methanol–water mixtures, 929
Matsuo, K.: see Tsuboi, M.
Mayer, A.: see Schlecht, P.; Casimir, W. v.
Merril, C. R.: Reinvestigation of the primary structure of yeast alanine tRNA, 1727
Merrill, E. W.: see Gibbs, D. A.
Miller, I. R., and Bach, D.: Interaction of DNA with heavy metal ions and polybases: Cooperative phenomena, 169
—: see Frommer, M. A.
Mitchell, T. W.: see Feughelman, M.
Miyazawa, T.: see Sugeta, H.
Montroll, E. W.: see Goel, N. S.
Morita, K., Simons, E. R., and Blout, E. R.: Polypeptides. LVI. Effect of lithium bromide and of temperature on the conformation of a copolymer of glutamic acid and lysine, 181

Nakahara, T.: see Itoh, K.
Nakajima, A., Hayashi, T., and Ohmori, M.: Phase equilibria of rodlike molecules in binary solvent systems, 973
Nakanishi, M.: see Tsuboi, M.
Noguchi, J.: see Kubota, S.

Oesper, P., and Scheinok, P. A.: On the discrimination between symmetric pentapeptides, 955
Ohmori, M.: see Nakajima, A.
Okabayashi, H., Isemura, T., and Sakakibara, S.: Steric structure of L-proline oligopeptides. II. Far-ultraviolet absorption spectra and optical rotations of L-proline peptides, 323
—: see Isemura, T.
Oosawa, F.: A theory on the effect of low molecular salts on the dissociation of linear polyacids, 135. A theory on the effect of low molecular salts on the conformation of linear polyions, 145. Interaction between parallel rodlike macroions, 1633

Oplatka, A., and Yonath, J.: Mechanochemical melting of collagen fibers. II, Diffusion-controlled contractions, 1147
—: see Reich, S.; Sherebrin, M. H.; Yonath, J.
Oya, M.: see Itoh, K.

Paleček, E., and Vetterl, V.: Polarographic reducibility of denatured DNA, 917
Pande, C. S., and McMenamy, R. H.: Association of thiocyanate with reduced bovine albumin: Essential features of the binding site, 1487
Pautard, F. G. E., and Zola, H.: Molecular weight of onuphic acid, 629
Peacocke, A. R., and Pritchard, N. J.: The ultrasonic degradation of biological macromolecules under conditions of stable cavitation. II. Degradation of deoxyribonucleic acid, 605
—: see Blake, A.
Peggion, E., Cosani, A., Verdini, A. S., Del Pra, A., and Mammi, M.: Conformational studies of poly-L-tryptophan: Circular dichroism and x-ray diffraction studies, 1477
—: see Cosani, A.; Goodman, M.
Pipkin, A. C.: see MacDonald, C. T.
Pritchard, N. J.: see Peacocke, A. R.
Profumo, P.: see Rialdi, G.
Puett, D., and Ciferri, A.: Pseudo-phase diagrams of poly-L-lysine and poly (L-glutamic acid) in aqueous salt solutions, 1213
—: see Ciferri, A.; Threlkeld, J. O.

Rajagh, L.: see Ciferri, A.
Ramachandran, G. N.: Need for nonplanar peptide units in polypeptide chains, 1494
—, and Chandrasekharan, R.: Interchain hydrogen bonds via bound water molecules in the collagen triple helix, 1649
—, and Venkatachalam, C. M.: Stereochemical criteria for polypeptides and proteins. IV. Standard dimensions for the *cis*-peptide unit and conformation of *cis*-polypeptides, 1255
—: see Sarathy, K. P.
Rattle, H. W. E.: see Bradbury, E. M.
Reich, S., Katchalsky, A., and Oplatka, A.: Dynamic-elastic investigation of the chemical denaturation of collagen fibers, 1159
—and Yonath, J.: Study of collagen denaturation kinetics by dynamic mechanical methods, 997
Rialdi, G., and Profumo, P.: Calorimetric measurements on DNA helix–coil transition in 2.2*M* urea, 899
Richter, M.: see Szejtli, J.
Ross, P. D., and Scruggs, R. L.: Viscosity study of DNA. II. The effect of simple salt concentration on the viscosity of high molecular weight DNA and application of viscometry to the study of DNA isolated from T4 and T5 bacteriophage mutants, 1005
Rubenstein, I.: see Gray, D. M.
Rüst, P.: Cesium chloride density distributions in preparative equilibrium ultracentrifugation, 1185

Sakakibara, S.: see Isemura, T.
Sarathy, K. P., and Ramachandran, G. N.: Possible formation of the left-handed α-helix of poly-L-serine, 461
Scheinok, P. A.: see Oesper, P.
Scheraga, H. A.: see Epand, R. F.; Ingwall, R. T.; Vournakis, J. N.
Schlecht, P., Vogel, H., and Mayer, A.: Effect of oxygen binding on the dielectric properties of hemoglobin, 1717
Schmit, G. N.: see Kovacs, J.
Schuster, T. M.: see Jennings, B. R.
Schwarz, G.: General theoretical approach to the thermodynamic and kinetic properties of cooperative intramolecular transformations of linear biopolymers, 873; *errata*, 1787
—, and Seelig, J.: Kinetic properties and the electric field effect of the helix–coil transition of poly(γ-benzyl L-glutamate) determined from dielectric relaxation measurements, 1263
Scott, R. A., III: Conformational studies of polynucleotides. I. Unperturbed dimensions in the approximation of independent units, 625
Scruggs, R. L.: see Ross, P. D.
Seelig, J.: see Schwarz, G.
Sharp, P., and Bloomfield, V. A.: Light scattering from wormlike chains with excluded volume effects, 1201
Sherebrin, M. H., and Oplatka, A.:

Contraction–relaxation of collagen fibers in LiBr–water–acetone solutions, 1169
Shimanouchi, T.: see Itoh, K.; Koyama, Y.
Shimshak, R. R.: see Gabbay, E. J.
Silberberg, A., and Simha, R.: Kinetics of reversible reactions on linear lattices with neighbor effects, 479
Simha, R.: see Silberberg, A.
Simons, E. R.: see Morita, K.
Sinsheimer, R. L.: see Strauss, J. H., Jr.
Smith, K. A.: see Gibbs, D. A.
Snaith, J. W.: see Haly, A. R.
Snart, R. S.: Effect of carcinogens and related compounds on the semiconductive properties of proteins, 73. Photoelectric effects of deoxyribonucleic acid, 293
Spach, G., and Constantin, D.: Poly-α-acide-L-glutamique en solution aqueuse. Agrégation spécifique et phénomènes d'hystérésis. II. Agrégation et précipitation. Analyse chromatographique, 653
—: see Jennings, B. R.
Stephens, R. M.: see Bradbury, E. M.
Sterling, C.: see Masuzawa, M.
Stewart, C. R.: Broadening by Acridine Orange of the thermal transition of DNA, 1737
Strauss, J. H., Jr., Kelly, R. B., and Sinsheimer, R. L.: Denaturation of RNA with dimethyl sulfoxide, 793
Sturtevant, J. M.: see Krakauer, H.
Sugai, S.: see Kubota, S.; Makino, S.
Sugeta, H., and Miyazawa, T.: A general method for calculating helical parameters of polymer chains from bond lengths, bond angles, and internal-rotation angles (*addendum*), 1387
Sundaralingam, M.: Some aspects of stereochemistry and hydrogen bonding of carbohydrates related to polysaccharide conformations, 189; *errata*, 1787
Suzuki, K., and Taniguchi, Y.: Effect of pressure on the helix and random coil forms of poly(D-glutamic acid), 215
Szejtli, J., Richter, M., and Augustat, S.: Molecular configuration of amylose and its complexes in aqueous solutions. IV. Determination of $\overline{DP}$ of amylose by measuring the concentration of free iodine in solution of amylose–iodine complex, 27; *errata*, 1787
Takashima, S.: Optical anisotropy of synthetic polynucleotides. I. Flow birefringence and π-electron polarizability of bases, 1437
Taniguchi, Y.: see Suzuki, K.
Terbojevich, M.: see Cosani, A.
Thompson, C. J.: Models for hemoglobin and allosteric enzymes, 1101
Threlkeld, J. O., Burke, J. J., Puett, D., and Cifferi, A.: Determination of the binding of tetraalkylammonium bromides to gelatin by an ultracentrifuge method, 767
Tiffany, M. L., and Krimm, S.: New chain conformations of poly(glutamic acid) and polylysine, 1379. Circular dichroism of poly-L-proline in an unordered conformation, 1767
Timasheff, S. N.: see Holcomb, D. N.
Tinoco, I., Jr.: see Davis, R. C.; Glaubiger, D.
Toniolo, C., Falxa, M. L., and Goodman, M.: Conformational aspects of polypeptides. XXV. Solvent and temperature effects on the conformations of copolymers of benzyl and methyl L-aspartate with nitrobenzyl L-aspartate, 1579
—: see Goodman, M.
Tooney, N. M., and Fasman, G. D.: Synthesis of poly-*O-tert*-butyl-L-serine and poly-L-serine, 81
Triebel, H.: Anomalies in the sedimentation behavior of high molecular weight DNA isolated from *Proteus mirabilis*, 449
Tsuboi, M., Matsuo, K., and Nakanishi, M.: Helix-with-loops structure of polynucleotide. II. Poly(I + CU), 123
Tunis, M.-J. B., and Hearst, J. E.: Optical rotatory dispersion of DNA in concentrated salt solutions, 1218. On the hydration of DNA. I. Preferential hydration and stability of DNA in concentrated trifluoroacetate solution, 1325. II. Base composition dependence of the net hydration of DNA, 1345
Turolla, A.: see Giacometti, G.

Uno, K.: see Itoh, K.

Venkatachalam, C. M.: Stereochemical criteria for polypeptides and proteins. V. Conformation of a system of three linked peptide units, 1425

—: see Ramachandran, G. N.
Venner, H.: see Zimmer, C.
Verdini, A. S.: see Cosani, A.; Peggion, E.
Vetterl, V.: see Paleček, E.
Vogel, H.: see Casimir, W. v.; Schlecht, P.
Vournakis, J. N., Yan, J. F., and Scheraga, H. A.: Effect of side chains on the conformational energy and rotational strength of the n–π^* transition for some α-helical poly-α-amino acids, 1531

Wakabayashi, K.: see Makino, S.
Walton, A. G.: see Carr, S. H.
Watanabe, H.: see Matsumoto, M.

Yamada, T., and Fukutome, H.: Vacuum ultraviolet absorption spectra of sublimed films of nucleic acid bases, 43
Yan, J. F.: see Vournakis, J. N.
Yang, J. T.: see Eyring, E. J.
Yew, F. F. H., and Davidson, N.: Breakage by hydrodynamic shear of the bonds between cohered ends of λ-DNA molecules, 659
Yonath, J., and Oplatka, A.: Mechanochemical melting of collagen fibers. I. Mechanical contractions, 1129
—: see Oplatka, A.; Reich, S.
Yoshioka, K.: see Matsumoto, M.

Zand, R.: Solution properties and structure of brain proteolipids, 939
Zaslow, B.: see Jackobs, J. J.
Zikán, J., and Kotýnek, O.: Interaction of antibodies specific towards the 2,4-dinitrophenyl group and of their fragments with hapten, 681
Zimm, B. H.: see Harpst, J. A.
Zimmer, C., Luck, G., Venner, H., and Frič, J.: Studies on the conformation of protonated DNA, 563
Zobel, H. F.: see Hinkle, M. E.
Zola, H.: see Pautard, F. G. E.

SUBJECT INDEX, VOLUME 6

Absorption spectra, far-ultraviolet, 323
  infrared, 307
  vacuum ultraviolet, 43
Acid–base behavior of copolymers, 1405
Acridine Orange, thermal transition broadening by, 1727
Acridine Orange–chondroitin sulfate C complexes, 703
Acridine–polynucleotide complexes, 1279
Adsorption of DNA, 1697
Aggregation, 653
  and precipitation, 653
  specific, of poly(α-L-glutamic acid), 635, 653
  temperature-dependent, 635
Albumin, reduced bovine, 1487
Alkali halide single crystals, 469
Amino acid derivatives, 993
Amino acids, 817
Aminoacridines, interaction with nucleic acids, 1225
9-Aminoacridine–DNA complex, 1279
Amylose, $\overline{DP}$ determination of, 27
  and its complexes, 27
Amylose–iodine complex, 27
Amylose–KBr, crystalline structure in oriented fibers of, 1659
Antianthrax immune serum, 817
Antibodies, specific toward 2,4-dinitrophenyl group, 681
Aromatic polypeptides, 1691
Aromatic side-chain effects on conformation, 1673
L-Aspartate esters, 837, 905
Assymetric surface, 993

Base compensation dependence of net hydration of DNA, 1345
Bases, flow birefringence and π-electron polarizability of, 1437
  nucleic acid, 43
Benzyl L-aspartate, 1579
Binding, of purine to polyuridylic acid, 983
  of tetraalkylammonium bromides to gelatin, 767
Binding isotherms, calculation of, 575
Binding sites, thiocyanate–albumin, 1487
Birefringence, flow, 105
  transient electric, 415
Bond angles, 1387
Bond lengths, 1387
Bonds, between cohered ends of λ-DNA molecules, 659
  hydrogen, 1649
Bound water in biopolymers, 1507
Bound water molecules in collagen triple helix, 1649
Branching, statistics of, 715

Calculating helical parameters of polymer chains, 1387
Calculations, for helix–coil transitions in heterogeneous polymers, 863
  of binding isotherms for heterogeneous polymers, 575
  of kinetic curves for the helix–coil transition of polypeptides, 385
  of melting curves for DNA, 1391
Calorimetric measurements on DNA helix–coil transitions, 899
Carbohydrates related to polysaccharide conformations, 189
Carboxymethyl cellulose, polyelectrolyte behavior of, 1519
Carcinogens and related compounds, 73
Cavitation, stable, conditions of, 605
Cellulose structure models, 771
Chain conformation, 1379
Chains, polymer, 1387
  polypeptide, 1494
Chondroitin sulfate C, 691
Chondroitin sulfate C–Acridine Orange complexes, 703
Chromatographic analysis, of aggregation and precipitation, 653
Circular dichroism, 1767
  of poly-L-tryptophan, 1477
Cis–trans equilibrium, 117
Collagen, denaturation kinetics by dynamic mechanical methods, 997
  triple helix, 1649
Complexes, chondroitin sulfate C–Acridine Orange, 703
  Poly A–poly U, 491
  2:1, between polyuridylic and oligoadenylic acids, 1737

Concentration of protein denaturants, 164
Configuration, molecular, of amylose and its complexes in aqueous solutions, 27
Conformation, aromatic side chain effects, 1673
of a copolymer of glutamic acid and lysine, 181
of linear polyions, 145
new chain, 1379
of poly-*O*-acetoallothreonine, 1311
of poly-*O*-acetylthreonine, 1311
of poly-L-methionone *S*-methylsulfonium salts in added salt systems, 551
of *cis*-polypeptides, 1255
of poly-L-tryptophan, 1477
of protonated DNA, 563
of a system of three linked peptide units, 1425
unordered, of poly-L-proline, 1767
Conformational aspects of polypeptides, 1579, 1691
Conformational assignments by far-UV Cotton effects, 1691
Conformational dependence of fluorescence polarization spectra, 159
Conformational energy of n–π^* transition, 1531
Conformational studies, of poly-L-alanine in water, 331
of polymers and copolymers of L-aspartate esters, 837, 905
of polynucleotides 625
of proteins, 1673
Conformations, polysaccharide, 189
of poly-L-valine in solution, 1551
Cooperative intramolecular transformations of linear biopolymers, 873
Cooperative phenomena, 169
Copoly(L-glutamic acid^{42}, L-lysine HBr^{28}, L-alanine^{30}), 851
Copolymer, dAT, 715
of glutamic acid and lysine, 181
of L-aspartate esters, 837
of benzyl and methyl L-aspartate with nitrobenzyl L-aspartate, 1579
synthetic periodic, 731
Copolypeptides in L-tryptophan, 159
Cotton effects, far-UV, 1691
Crystalline structure, in KBr–amylose, 1659
in oriented fibers, 1659
Crystallization, epitaxial, 469
Crystals, alkali halide single, 469
dAT, copolymer, 715
polymeric, 269
Decay processes in polydispersed systems 1491
Degradation, of deoxyribonucleic acid, 605
ultrasonic, 605
Denaturation, of DNA, 55, 751
of RNA with dimethyl sulfoxide, 793
Denaturation kinetics, collagen, 997
Denaturants, protein, 164
Denatured DNA, 917
Deoxyhemoglobin, 1705
Deoxyribonucleic acid, 293; see DNA.
degradation of, 605
solutions, 531
Dichloroacetic acid–1,2-dichloroethane mixtures, 1753
1,2-Dichloroethane–dichloroacetic acid mixtures, 1753
Dichroic ratio, ultraviolet, 1605
Dichroism, flow, 531
Dielectric dispersion, 1453
Dielectric properties, of hemoglobin, 1705
of oxyhemoglobin and deoxyhemoglobin, 1705
Dielectric relaxation measurements, 1263
Dimethyl sulfoxide, 793
2,4-Dinitrophenyl group, 681
Dinucleoside phosphates, 223, 409
DNA, adsorption at mercury–electrolyte interface, 1697
bacteriophage, 105, 595, 1605
calf thymus, 595, 1461
calorimetric measurements on, 899
denaturation and renaturation of, 731
denatured, 917
heat-denatured HeLa cell, 1573
helix–coil transition, 899
high molecular weight, 449
hydration of, 1325, 1345
influence of tritium labeling on physical properties of, 1461
interaction with heavy metals, 169
melting curves, 1391
net hydration of, 1345
optical properties of, 1509
preferential hydration of, 1325
protonated, 563
relaxation kinetics of denaturation of, 55
replicating and crosslinked, 285
solutions, 585
stability of in concentrated trifluoroacetate solutions, 1325

thermal transition of, 1727
ultraviolet dichroic ratios, 1605
DNA–9-aminoacridine complex, 1279
λ DNA, linear and cyclic, 1297
molecules, 659
$\overline{DP}$ of amylose, 27
DTA, detection and enthalpy of vaporization by, 1507
Dynamic mechanical methods, 997

Electrical properties of polypeptides in solution, 929
Electric birefringence in polydispersed systems, 1491
Electric field effect of helix–coil transition, 1263
π-Electron polarizability of bases, 1437
Electrophoresis of absorbed RNA, 97
Energy transfer, radiationless, 809
Enthalpy of helix–coil transitions from heats of solution, 441
of vaporization by DTA, 1507
Enthalpy change of helix–coil transition, 1745
Epitaxial crystallization of poly (γ-benzyl L-glutamate) on alkali halide single crystals, 469
Equilibria, hydrogen ion, 513
Esterolytic activity of vinylimidazole–vinylpyrrolidinone copolymers, 1417
Ethylene glycol, 1509

Far-infrared spectra, 1759
Far-ultraviolet absorption spectra, 323
Far-UV Cotton effects, 1691
Far ultraviolet optical rotatory properties of poly-L-tryptophan, 963
Films, sublimed, of nucleic acid bases, 43
Flow birefringence, of T2 bacteriophage DNA, 105
of bases, 1437
Flow dichroism of deoxyribonucleic acid solutions, 531
Formation of ω-helix, 905
β-Form structures, 1759
Free iodine, 27

Gelatin, 767
Gel filtration separation, theory of 1497
Gel–water relationships, 1453
Glutamic acid, 181

Hapten, 681
Heats of the helix–coil transitions of the poly A–poly U complexes, 491
Heats of solution, enthalpy of helix–coil transitions from, 441
Heavy metal ions, 169
HeLa cell DNA, heat-denatured, 1573
Helical parameters of polymer chains, calculating, 1386
α-Helical poly-α-amino acids, 1531
α-Helical structures, 1759
Helices, hairpin, of dAT copolymer, 715
nucleic acid, in solution, 993
Helix, collagen triple, 1649
ω-Helix, factors in formation of, 905
α-Helix, left-handed, 461
Helix and random coil forms of poly-(D-glutamic acid), 213
Helix–coil transitions, 863, 899, 1263
enthalpy of, from heats of solution, 441
enthalpy change of, 1745
heats of, 491
in heterogeneous polymers, 863
of poly(γ-benzyl L-glutamate), 1263
of poly-L-lysine, 1383
of polypeptides, 385
Helix-with-loops structure, of polynucleotide, 123
Hemoglobin, 1705
Heterogeneous polymers, calculation of binding isotherms for, 575
helix–coil transitions in, 863
High molecular weight DNA isolated from *Proteus mirabilis*, 449
Hyaluronate, polyelectrolyte behavior of, 1519
Hyaluronic acid, 777
Hydration, of DNA, 1325, 1345
of polyacids, 431
Hydrodynamic shear, 659
Hydrodynamical properties of polypeptides in solution, 929
Hydrogen bonding, C—H···O—C, 401
and stereochemistry of carbohydrates, 189
Hydrogen bonds, in bound water molecules, 1649
interchain, 1649
Hydrogen ion equilibria in poly(riboadenylic acid), 513
Hydrolysis, 255
RNAse catalyzed, 993
Hysteresis effects, in aqueous solutions, 635, 653

Infrared absorption spectra, 307, 323
Infrared spectra, low temperature, 401
Infrared studies of L-aspartate esters, 905

Interchain hydrogen bonds, 1649
Internal rotation angles, 1387
Intrinsic viscosities, of cyclic λ DNA, 1297
of linear λ DNA, 1297
Iodine–amylose complex, 27
Ion-binding of poly-L-methionone *S*-methylsulfonium salts in added salt systems, 551
Ionic polysaccharides, 1519
Ising model, one-dimensional, 117
Isotherms, binding, 575

KBr–amylose, crystalline structure in oriented fibers of, 1659
β-Keratin, 1515
Keratin fibers, 1503
thermally denatured 1515
Kinetics, of biopolymerization, 1
denaturation, 997
of primer-dependent polynucleotide phosphorylase synthetic reactions, 369
of reversible reactions on linear lattices 479
of synthesis of polymeric dAT, 269
Kinetic curves, calculation of, 385
Kinetic properties, of cooperative intramolecular transformations of linear biopolymers, 873
of helix–coil transition, 1263

Lattices, linear, 479
Light scattering, low-angle, 595
Light-scattering instrument, low-angle, 585
Linear biopolymers, 873
Linked peptide units, 1425
Lithium bromide, 181
Low-angle light scattering, 595
Low-angle light scattering instrument, 585
Low molecular salts, 135, 145
Low temperature infrared spectra, 401
Lysine, 181

Macroions, interaction between parallel rodlike, 1633
Macromolecules, biological, ultrasonic degradation of, 605
Mass spectrometry, thermal degradation of proteins studied by, 1001
Melting curves, DNA, 1391
Methanol–water, mixtures, 929
solvent systems, 1383
Methyl L-aspartate, 1579
Microwave frequencies, 1705
Mimimization approach for the improvement of cellulose structure models, 771
Model and parameters, possible use of synthetic periodic copolymers to establish, 731
Models, cellulose structure, 771
Molecular configuration of amylose, and its complexes in aqueous solutions, 27
Molecular conformation of poly(S-lactic acid), 299
Molecular weight, of onuphic acid, 629
of T7 and calf thymus DNA by low-angle light scattering, 595
Molecules, rodlike, 973

Neighbor effects, 479
Net hydration of DNA, 1345
Nitrobenzyl L-aspartate, 1579
Nucleic acid bases, sublimed films of, 43
Nucleic acid helices, 243, 255
in solution, 993
Nucleic acid templates, 1
Nucleic acids, interaction with aminoacridines, 1225

Oligoadenylic acid, 1737
Oligomeric templates, 269
Oligonucleotide interactions, 1737
Oligopeptides, 307, 323
One-dimenlional Ising model, 117
Onuphic acid, 629
Optical activity, of chondroitin sulfate C, 691
induced, 703
Optical anisotropy of synthetic polynucleotides, 1437
Optical properties, of DNA in ethylene glycol, 1509
temperature-dependent, 409
Optical rotation, of PGA, 635
of L-proline oligopeptides, 323
Optical rotatory properties, of poly-*O*-acetylallothreonine, 1311
of poly-*O*-acetylthreonine, 1311
of poly-L-tryptophan, 963
Optical spectroscopic studies of water-soluble polypeptides, 851
Oxygen binding, effect on hemoglobin, 1705
Oxyhemoglobin, 1705

Pentapeptides, 955
Peptide units, nonplanar, 1494
three linked, 1424

cis-Peptide unit, standard dimensions, 1255
Phase equilibria in rodlike molecules in binary solvent systems, 973
Phosphorylase, polynucleotide, 369
Photoelectric effects of deoxyribonucleic acid, 293
Polarization spectra, fluorescence, 159
Polarographic reducibility of denatured DNA, 917
Poly A–poly U complexes, 491
Poly-*O*-acetylallothreonine, 1311
Poly-*O*-acetylthreonine, 1311
Polyacids, hydration of, 431
  linear, 135
Polyadenylic acid, 255, 993
Polyadenylic–polyuridylic acid helices, 243
Poly-L-alanine, 331
Polyalanines with α-helical and β-form structures, 1759
Poly-α-amino acids, α-helical, 1531
Polyamines, 255
Polybases, 169
Poly(γ-benzyl L-glutamate), 469, 1263, 1745
Poly-*O*-*tert*-butyl-L-serine, 81
Polycytidylic–polyinosinic acid helices, 243
Polydispersed systems, 1491
Polyelectrolyte behavior of hyaluronate and carboxymethyl cellulose, 1519
Polyelectrolytes, limiting sedimentation coefficients of, 1500
Polyglutamates, 441
Poly(glutamic acid), 1379
Poly(D-glutamic acid), 213
Poly(L-glutamic acid), 851
  in methanol–water mixtures, 929
Poly(α-L-glutamic acid), specific aggregation of, 635, 653
Poly-γ-D- and L-glutamylglycine, 817
Polyglycine II, C—H···O—C hydrogen bonding in, 401
Polyglycines, low temperature infrared spectra of, 401
Poly(I + CU), 123
Polyinosinic–polycytidylic acid helices, 243
Polyions, linear, 145
Poly(S-lactic acid), 299
Polylysine, 1379
Poly-L-lysine, 1383
Poly-L-lysine HBr, 851
Polymer chains, 1387
Polymers, of L-aspartate esters, 837, 905
  heterogeneous, 575, 863
Poly-L-methionone *S*-methylsulfonium salts in added salt systems, 551
Polynucleotide–acridine complexes, 1279
Polynucleotide phosphorylase, 369
Polynucleotides, 123
  conformational studies of, 625
  synthetic, 1437
Polypeptide chain, 809, 1491
  in proteins, 164
Polypeptides, 181, 1255, 1425
  aromatic, 1691
  conformational aspects of, 1579, 1691
  electrical and hydrodynamical properties, 929
  helix–coil transition of, 385
  with known repeating sequences of amino acids, 817
  water-soluble, 851
cis-Polypeptides, conformation of, 1255
Poly-L-proline, 117, 307
  in unordered conformation, 1767
Poly(riboadenylic acid), 513
Polysaccharide conformations, 189
Polysaccharides, ionic, 1519
  synthetic and natural, 1573
Poly-L-serine, 81, 461
Poly-L-tryptophan, 963, 1477
Poly U–poly A complexes, 491
Polyuridylic acid, 1737
  binding of purine to, 983
Polyuridylic–polyadenylic acid helices, 243
Poly-L-valine, conformations of in solution, 1551
Poly(vinylpyrrolidone–vinylimidazole) copolymers, 1405
Pore sizes, 1497
Precipitation, 653
  and aggregation, 653
Preferential hydration of DNA, 1325
Preparation of L-aspartate esters, 837
Pressure, effect of, 213
Primary structure of tRNA, 1717
Primer-dependent synthetic reactions, 369
L-Proline, 307, 323
L-Proline oligopeptides, 323
Protein denaturants, 164
Proteins, 164, 1255, 1425
  conformation of, 1673
  semiconductive properties of, 73
  thermal degradation of, 1001
Proteolipids, brain, 939

Interchain hydrogen bonds, 1649
Internal rotation angles, 1387
Intrinsic viscosities, of cyclic λ DNA, 1297
of linear λ DNA, 1297
Iodine–amylose complex, 27
Ion-binding of poly-L-methionone *S*-methylsulfonium salts in added salt systems, 551
Ionic polysaccharides, 1519
Ising model, one-dimensional, 117
Isotherms, binding, 575

KBr–amylose, crystalline structure in oriented fibers of, 1659
β-Keratin, 1515
Keratin fibers, 1503
thermally denatured 1515
Kinetics, of biopolymerization, 1
denaturation, 997
of primer-dependent polynucleotide phosphorylase synthetic reactions, 369
of reversible reactions on linear lattices 479
of synthesis of polymeric dAT, 269
Kinetic curves, calculation of, 385
Kinetic properties, of cooperative intramolecular transformations of linear biopolymers, 873
of helix–coil transition, 1263

Lattices, linear, 479
Light scattering, low-angle, 595
Light-scattering instrument, low-angle, 585
Linear biopolymers, 873
Linked peptide units, 1425
Lithium bromide, 181
Low-angle light scattering, 595
Low-angle light scattering instrument, 585
Low molecular salts, 135, 145
Low temperature infrared spectra, 401
Lysine, 181

Macroions, interaction between parallel rodlike, 1633
Macromolecules, biological, ultrasonic degradation of, 605
Mass spectrometry, thermal degradation of proteins studied by, 1001
Melting curves, DNA, 1391
Methanol–water, mixtures, 929
solvent systems, 1383
Methyl L-aspartate, 1579
Microwave frequencies, 1705
Mimimization approach for the improvement of cellulose structure models, 771
Model and parameters, possible use of synthetic periodic copolymers to establish, 731
Models, cellulose structure, 771
Molecular configuration of amylose, and its complexes in aqueous solutions, 27
Molecular conformation of poly(S-lactic acid), 299
Molecular weight, of onuphic acid, 629
of T7 and calf thymus DNA by low-angle light scattering, 595
Molecules, rodlike, 973

Neighbor effects, 479
Net hydration of DNA, 1345
Nitrobenzyl L-aspartate, 1579
Nucleic acid bases, sublimed films of, 43
Nucleic acid helices, 243, 255
in solution, 993
Nucleic acid templates, 1
Nucleic acids, interaction with aminoacridines, 1225

Oligoadenylic acid, 1737
Oligomeric templates, 269
Oligonucleotide interactions, 1737
Oligopeptides, 307, 323
One-dimenlional Ising model, 117
Onuphic acid, 629
Optical activity, of chondroitin sulfate C, 691
induced, 703
Optical anisotropy of synthetic polynucleotides, 1437
Optical properties, of DNA in ethylene glycol, 1509
temperature-dependent, 409
Optical rotation, of PGA, 635
of L-proline oligopeptides, 323
Optical rotatory properties, of poly-*O*-acetylallothreonine, 1311
of poly-*O*-acetylthreonine, 1311
of poly-L-tryptophan, 963
Optical spectroscopic studies of water-soluble polypeptides, 851
Oxygen binding, effect on hemoglobin, 1705
Oxyhemoglobin, 1705

Pentapeptides, 955
Peptide units, nonplanar, 1494
three linked, 1424

cis-Peptide unit, standard dimensions, 1255
Phase equilibria in rodlike molecules in binary solvent systems, 973
Phosphorylase, polynucleotide, 369
Photoelectric effects of deoxyribonucleic acid, 293
Polarization spectra, fluorescence, 159
Polarographic reducibility of denatured DNA, 917
Poly A–poly U complexes, 491
Poly-*O*-acetylallothreonine, 1311
Poly-*O*-acetylthreonine, 1311
Polyacids, hydration of, 431
  linear, 135
Polyadenylic acid, 255, 993
Polyadenylic–polyuridylic acid helices, 243
Poly-L-alanine, 331
Polyalanines with α-helical and β-form structures, 1759
Poly-α-amino acids, α-helical, 1531
Polyamines, 255
Polybases, 169
Poly(γ-benzyl L-glutamate), 469, 1263, 1745
Poly-*O*-*tert*-butyl-L-serine, 81
Polycytidylic–polyinosinic acid helices, 243
Polydispersed systems, 1491
Polyelectrolyte behavior of hyaluronate and carboxymethyl cellulose, 1519
Polyelectrolytes, limiting sedimentation coefficients of, 1500
Polyglutamates, 441
Poly(glutamic acid), 1379
Poly(D-glutamic acid), 213
Poly(L-glutamic acid), 851
  in methanol–water mixtures, 929
Poly(α-L-glutamic acid), specific aggregation of, 635, 653
Poly-γ-D- and L-glutamylglycine, 817
Polyglycine II, C—H···O—C hydrogen bonding in, 401
Polyglycines, low temperature infrared spectra of, 401
Poly(I + CU), 123
Polyinosinic–polycytidylic acid helices, 243
Polyions, linear, 145
Poly(S-lactic acid), 299
Polylysine, 1379
Poly-L-lysine, 1383
Poly-L-lysine HBr, 851
Polymer chains, 1387
Polymers, of L-aspartate esters, 837, 905
  heterogeneous, 575, 863
Poly-L-methionone *S*-methylsulfonium salts in added salt systems, 551
Polynucleotide–acridine complexes, 1279
Polynucleotide phosphorylase, 369
Polynucleotides, 123
  conformational studies of, 625
  synthetic, 1437
Polypeptide chain, 809, 1491
  in proteins, 164
Polypeptides, 181, 1255, 1425
  aromatic, 1691
  conformational aspects of, 1579, 1691
  electrical and hydrodynamical properties, 929
  helix–coil transition of, 385
  with known repeating sequences of amino acids, 817
  water-soluble, 851
cis-Polypeptides, conformation of, 1255
Poly-L-proline, 117, 307
  in unordered conformation, 1767
Poly(riboadenylic acid), 513
Polysaccharide conformations, 189
Polysaccharides, ionic, 1519
  synthetic and natural, 1573
Poly-L-serine, 81, 461
Poly-L-tryptophan, 963, 1477
Poly U–poly A complexes, 491
Polyuridylic acid, 1737
  binding of purine to, 983
Polyuridylic–polyadenylic acid helices, 243
Poly-L-valine, conformations of in solution, 1551
Poly(vinylpyrrolidone–vinylimidazole) copolymers, 1405
Pore sizes, 1497
Precipitation, 653
  and aggregation, 653
Preferential hydration of DNA, 1325
Preparation of L-aspartate esters, 837
Pressure, effect of, 213
Primary structure of tRNA, 1717
Primer-dependent synthetic reactions, 369
L-Proline, 307, 323
L-Proline oligopeptides, 323
Protein denaturants, 164
Proteins, 164, 1255, 1425
  conformation of, 1673
  semiconductive properties of, 73
  thermal degradation of, 1001
Proteolipids, brain, 939

Proton magnetic resonance study, of the binding of purine to polyuridylic acid, 983
of water-soluble polypeptides, 851
Purine, binding of to polyuridylic acid, 983

Radiationless energy transfer through a polypeptide chain, 809
Random coil and helix forms of poly(D-glutamic acid), 213
Relaxation kinetics of denaturation of DNA, 55
Renaturation of DNA, 731
Rheology of hyaluronic acid, 777
RNA, absorbed, 97
denaturation of, 793
RNAse-catalyzed hydrolysis, 993
of polyadenylic acid, 255
tRNA, yeast alanine, 1717
Rodlike macroions, 1633
Rodlike molecules, phase equilibria in, 973
Rotational strength of n–π^* transition, 1531

Salts, low molecular, 135, 145
poly-L-methionone *S*-methylsulfonium, 551
Sedimentation behavior, anomalies in, 449
Sedimentation coefficients, of replicating and crosslinked DNA, 285
limiting, 1500
Semiconductive properties of proteins, 73
Serological reaction, 817
Shear, hydrodynamic, 659
Side-chain effects, aromatic, 1673
on n–π^* transition, 1531
Solution properties and structure of brain proteolipids, 939
Solution studies of L-aspartate esters, 837
Solvent effects on conformations, 1579
Solvent systems, binary, 973
Specific heat studies, 1355
Spectra, far-infrared, 1759
far-ultraviolet absorption, 323
fluorescence polarization, 159
infrared absorption, 307
low temperature infrared, 401
vacuum ultraviolet absorption, 43
Spermine and spermidine derivatives, 243
Stability of DNA in concentrated trifluouroacetate solution, 1325
Statistics of branching and hairpin helices for the dAT copolymer, 715
Strain–stress isotherm of keratin fibers, 1503
Structure, of brain proteolipids, 939
of tRNA, 1717
Stereochemical criteria for polypeptides and proteins, 1255, 1425
Stereochemistry and hydrogen bonding of carbohydrates, 189
Steric structure of L-proline oligopeptides, 307, 323
Symmetric pentapeptides, discrimination between, 955
Synthesis, kinetics of, 269
of poly-*O-tert*-butyl-L-serine and poly-L-serine, 81
of poly-γ-D- and L-glutamylglycine, 817
Synthetic polynucleotides, 1437
Synthetic reactions, polynucleotide phosphorylase, 369
Systems, added salt, 551
Sublimed films of nucleic acid bases, 43

Temperature, effect of on conformation, 181, 1579
Temperature dependence of the hydrogen ion equilibria in poly(riboadenylic acid), 513
Temperature-dependent aggregation, 635
Temperature-dependent optical properties of a torsional oscillator model for dinucleoside phosphates, 409
Temperature-dependent properties of dinucleoside phosphates, 223
Templates, nucleic acid, 1
oligomeric, 269
Tetraalkylammonium bromides, 767
Thermal degradation of proteins, 1001
Thermal transition of DNA, 1727
Thermally denatured keratin fibers, 1515
Thermodynamic properties of cooperative intramolecular transformations of linear biopolymers, 873
Thiocyanate, 1487
Topography of nucleic acid helices in solution, 243, 255, 993
Torsional oscillator model for dinucleoside phosphates, 409
Transient electric birefringence studies of T2 bacteriophage and T2 ghost, 415
n–π^* Transition, effect of side chains on conformational energy and retational strength of, 1531
Trifluoroacetate solution, concentrated, 1325

Triple helical single-chain structures, 1771
Triplet energy delocalization, 1279
Triplet states, 1279
Tritium labeling, 1461
L-Tryptophan containing copolypeptides, 159

Ultracentrifuge method, 767
Ultraviolet absorption spectra, vacuum, 43
Ultraviolet dichroic ratio, 1605

Vacuum ultraviolet absorption spectra, 43
Vinylimidazole–vinylpyrrolidinone copolymers, 1417
Vinylpyrrolidinone–vinylimidazole copolymers, 1417
Viscosity of chondroitin sulfate C, 691
Volume fraction of the polypeptide chain in proteins, 164

Water, bound in bipolymers, 1507, 1649
Wilzbach method, 1461
Wool–water systems, 1355

X-Ray diffraction of poly-L-tryptophan, 1477